

C-01			
IUPAC	formula	amt	qnt
Ammonium persulfate, 98+% ; A.C.S. reagent	$(\text{NH}_4)_2\text{S}_2\text{O}_8$	100g	1

C-04			
IUPAC	formula	amt	qnt
2,6-Naphthalenedicarboxylic acid, 97%	$\text{C}_{12}\text{H}_8\text{O}_4$	5 g	1
L(+) – Ascorbic acid, 99%	$\text{C}_6\text{H}_8\text{O}_6$	500g	1
Acetic acid sodium salt trihydrate, p.a.	$\text{C}_2\text{H}_3\text{NaO}_2 \cdot 3\text{H}_2\text{O}$	1 KG	1

C-05			
IUPAC	formula	amt	qnt
4, 4' – Biphenol, 97%	$\text{C}_{12}\text{H}_{10}\text{O}_2$	100g	2
Phenolphthalein, 98.5% pure, Indicator grade	$\text{C}_{20}\text{H}_{14}\text{O}_4$	100g	2
2, 2, 3, 3, 4, 4, 5, 5, 6, 6, 7, 7, 8, 8, 9, 9 – Hexadecafluoro – 1, 10 – decanediol, 96%		5 g	1
4, 4' – Dihydroxybenzophenon	$(\text{HOC}_6\text{H}_4)_2\text{CO}$	25g	1
1, 1, 1 – Tris(4 – hydroxyphenyl)ethane, 99%	$\text{C}_{20}\text{H}_{18}\text{O}_3$	100g	1
2 – pentanone, 97%	$\text{C}_5\text{H}_{10}\text{O}$	500ml	1
2 – mercaptoethanol, 98%	$\text{C}_2\text{H}_6\text{OS}$	250ml	1
2 – mercaptoethanol, 99%	$\text{C}_2\text{H}_6\text{OS}$	250ml	1
Octyl ether, 99%		25g	1
1, 2 – Tetradecandiol, technical grade, 90%	$\text{CH}_3(\text{CH}_2)_{11}\text{CH}(\text{OH})\text{CH}_2\text{OH}$	100g	1
1, 4 – Butanediol, diglycidyl ether, 50~60%, tech	$\text{C}_{10}\text{H}_{18}\text{O}_4$	100g	1
4 – Aminophenethyl alcohol, 97%	$\text{C}_8\text{H}_{11}\text{NO}$	25g	1
4, 4' – (Hexafluoroisopropylidene)diphenol, 97%	$\text{C}_{15}\text{H}_{10}\text{F}_6\text{O}_2$	25g	1
1, 8 – Octanediol, 98+%		100g	1
Hydroquinone sulfonic acid, potassium salt, tech	$\text{C}_6\text{H}_5\text{KO}_5\text{S}$	100g	1
Phloroglucinol, 99+%, anhydrous	$\text{C}_6\text{H}_6\text{O}_3$	50g	1
Isopropanol, For analysis	$(\text{CH}_3)_2\text{CHOH}$	1 L	1
Oxalic acid, 98%, anhydrous	$\text{C}_2\text{H}_2\text{O}_4$	250g	1
1, 2 – Hexadecanediol, tech, 90%	$\text{C}_{16}\text{H}_{34}\text{O}_2$	50g	1
3, 3' - Dihydroxybenzidine	$\text{C}_{12}\text{H}_{12}\text{N}_2\text{O}_2$	25g	1
Chromotropic acid disodium salt dihydrate	$\text{C}_{10}\text{H}_6\text{Na}_2\text{O}_8\text{S}_2 \cdot 2\text{H}_2\text{O}$	25g	1
4 – Phenylphenol, 97%	$\text{C}_{12}\text{H}_{10}\text{O}$	250g	1
Resorcinol, 98%	$\text{C}_6\text{H}_6\text{O}_2$	250g	1
N – Hydroxysuccinimide, 98%	$\text{C}_4\text{H}_5\text{NO}_3$	25g	1
2 – Bromoethanol, 97%	$\text{C}_2\text{H}_5\text{BrO}$	100ml	1
7 – Hydroxyl – 4 – methylcoumarin, 97% pure	$\text{C}_{10}\text{H}_8\text{O}_3$	100g	1

N – Bromosuccinimide, 99%	C ₄ H ₄ BrNO ₂	500g	1
4, 4' - Thiodiphenol	C ₁₂ H ₁₀ O ₂ S	100g	1
Bisphenol A diglycidyl ether	C ₂₁ H ₂₄ O ₄	250g	1

C-06			
IUPAC	formula	amt	qnt
Tetraethylene glycol, 99.50%	C ₈ H ₁₈ O ₅	1 L	2
Polyethylene Glycol, 4000	HO(CH ₂ CH ₂ O) _n CH ₂ CH ₂ OH	500g	1
4 – tert – Butylphenol, 97%	C ₁₀ H ₁₄ O	1 KG	1
4 – Nitrophenol, 98%	C ₆ H ₅ NO ₃	1 KG	1
Benzyl alcohol, 99% pure	C ₇ H ₈ O	1 L	1
3, 3', 4, 4' – Benzophenonetetracarboxylic dianhydride, 99%	C ₁₇ H ₆ O ₇	500g	1
4, 4' – Sulfonyldiphenol, 99.70%	C ₁₂ H ₁₀ O ₄ S	500g	1
M - Nitrophenol	NO ₂ C ₆ H ₄ OH	500g	1
Hexyl alcohol, 98%	C ₆ H ₁₄ O	1 L	1

C-07			
IUPAC	formula	amt	qnt
N – Nonane, 99%	C ₉ H ₂₀	100ml	1
1 – Bromobutane, 99%	C ₄ H ₉ Br	1 L	1

D-09			
IUPAC	Formula	amt	qnt
Ethyl alcohol, 99.5%	C ₂ H ₅ OH	500ml	1